

结晶紫-柠檬酸染色液(0.1%)

货号: G1074

规格: 100mL

保存: 室温, 避光保存, 有效期1年。

产品介绍:

结晶紫是一种可以和细胞核中的 DNA结合的碱性染料, 对细胞核染色。结晶紫染色切片采用水溶性封固剂封固是很必要的, 因为局部的脱水作用会破坏红色的多染性。

结晶紫-柠檬酸染色液(0.1%)主要由结晶紫、柠檬酸等组成, 是一种组织或细胞染色时常用的可以把细胞核染成深紫色的染色液。

自备材料:

系列乙醇、蒸馏水、4%多聚甲醛

操作步骤: (仅供参考)

1. 样品处理:

石蜡切片: 二甲苯中脱蜡 2 次每次 5-10min。系列乙醇(100%、95%、85%、75%)复水, 每梯度 3min。蒸馏水 2min。

冰冻切片: 取出恢复至室温后蒸馏水浸洗 2min。

培养细胞: 用4%多聚甲醛固定10min以上。蒸馏水清洗2次, 每次2min。

2. 染色:

结晶紫-柠檬酸染色液(0.1%)染色10min(可以根据染色结果和要求调整时间)。

用蒸馏水或自来水充分洗涤, 可进行观察和拍照。

注意事项:

1. 如果需要脱水、透明和封片处理, 需自备二甲苯, 中性树胶或其它封片剂。
2. 第一次使用本试剂时建议先取 1~2个样品做预实验。
3. 为了您的安全和健康, 请穿实验服并戴一次性手套操作。

Crystal Violet - Citric Acid Stain Solution, 0.1%

Cat: G1074

Size: 100mL

Storage: RT, avoid light, valid for 1 year.

Introduction

Crystal violet is an alkaline dye that binds to DNA in the nucleus and stains the nucleus. It is necessary for methyl violet staining sections to be sealed with water-soluble cementing agents, because local dehydration will destroy the red polychromatism.

Crystal Violet - Citric Acid Stain Solution, 0.1% is a dyeing solution commonly used in tissue or cell dyeing, which can dye the nucleus into deep purple.

Self Provided Materials

Gradient ethanol, distilled water, 4% polyformaldehyde

Protocol(*for reference only*)

1. Sample treatment

For paraffin section: dewax in xylene twice for 5-10mins each. Rehydrate with series ethanol (100%,90%,80%,70%) for each level 3mins,and finally in distilled water for 3mins.

For frozen section: restore to RT and wash with distilled water for 2mins.

For cultured cell: fix with 4% PFA for more than 10mins.Then wash with distilled water twice for 2mins each.

2. Crystal violet stain

- 1) Crystal Violet - Citric Acid Stain Solution, 0.1% dyes for 10 min (the dyeing time could be adjusted according to the dyeing results and requirements).
- 2) View and photograph after fully washing with distilled water or tap water.

Note

1. If dehydration, transparency and sealing are needed, self-prepared xylene, neutral gum or other sealing agents are required.
2. It is suggested to take 1-2 samples for preliminary experiment to determine the dilution ratio and dyeing time of the dye solution before formal experiment.
3. For your safety and health, please wear experimental clothes and disposable gloves.