

TECHNICAL DATA SHEET

violetFluor™ 450 Anti-Mouse Ly-6G/Ly-6C (Gr-1) (RB6-8C5)

Catalog Number: 75-5931

PRODUCT INFORMATION

Contents: violetFluor™ 450 Anti-Mouse Ly-6G/Ly-6C (Gr-1)

Isotype: Rat IgG2b, kappa

Concentration: 0.2 mg/mL


Clone: RB6-8C5

Reactivity: Mouse

Use By: 6 months from date of receipt

Storage Conditions: 2-8°C protected from light

Formulation: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% NaN₃, 0.1% gelatin, pH7.2



Mouse Ly-6G (RB6-8C5) violetFluor™ 450

C57Bl/6 bone marrow cells were stained with 0.06 ug violetFluor™ 450 Anti-Mouse Ly-6G/Ly-6C (75-5931) (solid line) or 0.06 ug violetFluor™ 450 Rat IgG2b isotype control (dashed line).

DESCRIPTION

The RB6-8C5 antibody binds to mouse Ly-6G/Ly-6C, commonly known as Gr-1, a member of the Ly-6 superfamily of GPI-anchored cell surface proteins with roles in cell signaling and cell adhesion. Gr-1 is differentially expressed during development and maturation of cells in the myeloid lineage and is expressed at varying stages and levels on monocytes, macrophages, granulocytes, and peripheral neutrophils.

PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

APPLICATION NOTES

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). Please refer to the figure legend for the optimal concentration used to stain the tissue shown. We recommend titrating the antibody under your specific conditions to determine the optimal concentration of antibody needed in your experimental system.

REFERENCES

Berent-Maoz B, Montecino-Rodriguez E, Signer RAJ, and Dorshkind K. 2012. Blood. 199:5715-5721. (Flow cytometry) von Bruhl M-L, Stark K, Steinhart A, et al. 2012. J. Exp. Med. 209: 819-835. (Intravital fluorescent microscopy - video) Le HT, Tran VG, Kim W, Kim H, Cho HR, and Kwon B. 2012. J. Immunol. 189:287-295. (in vivo neutrophil depletion) Doring Y, Soehnlein O, Drechsler M, Shagdarsuren E, Chaudhari SM, Meiler S, Hartwig H, Hristov M, Koenen RR, Hieronymus T, Zenke M, Weber C, and Zernecke A. 2012. Arterioscler. Thromb. Vasc. Biol. 32: 1613-1623. (in vivo depletion) Hickman HD, Li L, Reynoso GV, Rubin EJ, Skon CN, Mays JW, Gibbs J, Schwartz O, Bennink JR, and Yewdell JW. 2011. J. Exp. Med. 208: 2511-2524. (Immunohistochemistry – OCT embedded frozen tissue) Wang T, Tian L, Haino M, Gao J-L, Lake R, Ward Y, Wang H, Siebenlist U, Murphy PM, and Kelly K. 2007. Infect. Immun. 75(3):1144-1153. (Immunohistochemistry – zinc fixed tissue) Nutt SL, Metcalf D, D'Amico A, Polli M, and Wu L. 2005. J. Exp. Med. 201:221-231. (Immunomagnetic bead depletion) Whiteland JL, Nicholls SM, Shimeld C, Easty DL, Williams NA, and Hill TJ. 1995. J. Histochem. Cytochem. 43:313-320. (Immunohistochemistry – frozen tissue, paraffin embedded tissue) Fleming TJ, Fleming ML, and Malek TR. 1993. J. Immunol. 151:2399-2408. (in vitro blocking, immunoprecipitation)

Cytek Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Cytek Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.