

GAPDH Rabbit mAb

Catalog :

TS30006

Gene Name :

GAPDH

Gene ID :

Human: 2597

Mouse:100042025

Rat:24383

SwissProt :

Human:P04406

Mouse:P16858

Rat:P04797

Host :

Rabbit

Clonality :

Monoclonal

Clone No. :

TSKL30006

Reactivity :

Human,Mouse,Rat

Storage :

-15°C to -25°C/1 year
(Do not lower than -25°C)

Specificity :

Endogenous

Purification :

Protein A

Molecular Weight :

MW(Calculated): 38 kDa

MW(Observed): 38 kDa

Application :

WB,IHC,IF,IP,ELISA

Dilution :


IHC 1:2000-1:5000

WB 1:5000-1:25000

IF 1:50-1:200

ELISA 1:5000-1:20000


IP 1:50-1:200



Immunohistochemical analysis of paraffin-embedded Human breast tumor tissue using GAPDH Rabbit mAb. The section was pre-treated using heat mediated antigen retrieval with Tris/EDTA buffer (pH9.0) for 20 minutes. The tissues were probed with the primary antibody (TS30006, 1/400) overnight at 4°C. The polymer-HRP anti-Rabbit IgG (H+L) antibody used to detect the antibody.

Western blot analysis of various lysates using GAPDH Rabbit mAb (TS30006) at 1:10000 dilution.

Image



Lane 1: HeLa cell lysate

Lane 2: MCF7 cell lysate

Lane 3: A549 cell lysate

Lysates/proteins at 20 µg/Lane

Predicted band size: 38 kDa

Observed band size: 38 kDa

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5% nonfat dry milk in TBST.

Exposure time: 5 s (pg-level)

Reactivity Key: H=Human; M=Mouse; R=Rat; Mk=Monkey; Hm=Hamster

Application Key: WB=Western Blot; IHC-P=Immunohistochemistry (Paraffin); ICC=Immunocytochemistry;

IF=Immunofluorescence; FC=Flow Cytometry; IP=Immunoprecipitation

01


GAPDH Rabbit mAb

Storage Buffer :

PBS,50%glycerol, 0.05%Proclin
300, 0.05%BSA

Western blot analysis of various lysates using GAPDH Rabbit mAb (TS30006) at 1:10000 dilution.

Image



Lane 1: Mouse brain tissue lysate

Lane 2: Rat brain tissue lysate

Lysates/proteins at 30 µg/Lane

Predicted band size: 38 kDa

Observed band size: 38 kDa

Secondary antibody:HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution.Blocking buffer: 5%nonfat dry milk in TBST.

Exposure time:2 s(pg-level)

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster

Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;

IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation

02