
Orders:market@tsingke.com.cn
Support:protein@tsingke.com.cn
Web:www.tsingke.com.cn

Histone H3 Rabbit mAb


Catalog :
TS30005
Gene Name :
HIST1H3A
Gene ID :
Human:8350;
8351;
8352;
8353;
8354;
8355;
8356;
8357;
8358;
8968;
Mouse:319152;
Rat:291159;
SwissProt :
Human:P68431
Mouse:P68433
Rat:Q6LED0
Host :
Rabbit
Clonality :
Monoclonal
CloneNo. :
TSKL30005
Reactivity :
Human,Mouse,Rat
Storage :
-15°C to -25°C/1 year
(Do not lower than -25°C)
Specificity :
Endogenous
Purification :
Protein A
Molecular Weight :
MW(Calculated): 15 kD
MW(Observed): 15 kD



Immunohistochemical analysis of paraffin- Histone H3 Rabbit mAb.The section was pre-treated using heat mediated antigen retrieval with Tris/EDTA buffer (pH9.0) for 20 minutes. The tissues were probed with the primary antibody (TS30005,1/400) overnight at 4℃.The polymer-HRP anti-Rabbit IgG(H+L) antibody used to detect the antibody.

Western blot analysis of various lysates using Histone H3 Rabbit mAb (TS30005)at 1:5000 dilution.

Image



Lane 1: HeLa cell lysate
Lane 2: HEK293 cell lysate

Lysates/proteins at 15μg/Lane

Predicted band size: 15 kDa
Observed band size: 15 kDa

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5%nonfat dry milk in TBST.

Exposure time:1 s(pg-level)

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster
Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;
IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation

For Research Use Only.Not for Use in Diagnostic Procedures.


Orders:market@tsingke.com.cn
Support:protein@tsingke.com.cn
Web:www.tsingke.com.cn

Histone H3 Rabbit mAb

Application :
WB,IHC,IF,IP,ELISA
Dilution :
IHC 1:5000-1:20000 WB 1:5000-1:10000
IF 1:200-1:1000
ELISA 1:5000-1:20000
IP 1:50-1:200
Storage Buffer :
PBS,50%glycerol, 0.05%Proclin 300, 0.05%BSA

Western blot analysis of various lysates using Histone H3 Rabbit mAb(TS30005)at 1:5000 dilution.

Image



Lane 1: Mouse liver tissue lysate
Lane 2: Rat liver tssue lysate
Lane 3: Mouse lung tissue lysate
Lane 4: Rat lung tissue lysate

Lysates/proteins at 30 μg/Lane

Predicted band size:15 kDa
Observed band size:15 kDa

Secondary antibody:HRP-conjugated Goat anti-Rabbit IgG(H+L)at 1:10000 dilution.Blocking buffer:5%nonfat dry milk in TBST.

Exposure time:1 s(pg-level)

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster
Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation

For Research Use Only.Not for Use in Diagnostic Procedures.