

p70 S6 Kinase(Phospho Thr389) Rabbit mAb

Catalog :

TS10082

Gene Name :

p70 S6 kinase α

Gene ID :

Human: 6198

Mouse: 72508

Rat: 83840

SwissProt :

Human: P23443

Mouse: Q8BSK8

Rat: P67999

Host :

Rabbit

Clonality :

Monoclonal

Clone No. :

TSKL10082

Reactivity :

Human,Mouse,Rat

Storage :

-15°C to -25°C/1 year
(Do not lower than -25°C)

Specificity :

Endogenous

Purification :

Protein A

Molecular Weight :

MW(Calculated): 59 kD

MW(Observed): 70 kD

Application :

WB,IP,ELISA

Dilution :

WB 1:1000-1:5000

ELISA 1:5000-1:20000


IP 1:50-1:200

Storage Buffer :

PBS,50%glycerol,0.05%Proclin
300,0.05%BSA

Western blot analysis of various lysates using p70 S6 Kinase (Phospho Thr389) Rabbit mAb (TS10082) at 1:2000 dilution.

Image



Lane 1: MCF7 cell lysate

Lane 2: C6 cell lysate

Lysates/proteins at 15 µg/Lane

Predicted band size: 59 kDa

Observed band size: 70 kDa

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG(H+L)
at1:10000 dilution. Blocking buffer: 5%nonfat dry milk in TBST.

Exposure time: 45 s(fg-level).

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster

Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;

IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation

01

p70 S6 Kinase(Phospho Thr389) Rabbit mAb

Catalog :

TS10082

Gene Name :

p70 S6 kinase α

Gene ID :

Human: 6198

Mouse: 72508

Rat: 83840

SwissProt :

Human: P23443

Mouse: Q8BSK8

Rat: P67999

Host :

Rabbit

Clonality :

Monoclonal

Clone No. :

TSKL10082

Reactivity :

Human,Mouse,Rat

Storage :

-15°C to -25°C/1 year
(Do not lower than -25°C)

Specificity :

Endogenous

Purification :

Protein A

Molecular Weight :

MW(Calculated): 59 kDa

MW(Observed): 70 kDa

Application :

WB,IP,ELISA

Dilution :

WB 1:1000-1:5000

ELISA 1:5000-1:20000


IP 1:50-1:200

Storage Buffer :

PBS,50%glycerol,0.05%Proclin
300,0.05%BSA

Western blot analysis of various lysates using p70 S6 Kinase (Phospho Thr389) Rabbit mAb (TS10082) at 1:2000 dilution.

Image



Lane 1: NIH-3T3 cell lysate

Lane 2: Overnight serum-starved,
Calyculin A-treated NIH-3T3 cell
lysate

Lane 3: Overnight serum-starved,
IGF-1-treated NIH-3T3 cell lysate

Lysates/proteins at 20 µg/Lane

Predicted band size: 59 kDa

Observed band size: 70 kDa

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5%BSA in TBST.

Exposure time: Lane 1-2:3 s(pg-level);Lane 3:37 s(fg-level).

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster

Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;
IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation