

p53(Phospho Ser392) Rabbit mAb

Catalog :

TS10088

Gene Name :

p53

Gene ID :

Human: 7157

Mouse: 22059

Rat: 24842

SwissProt :

Human: P04637

Mouse: P02340

Rat: P10361

Host :

Rabbit

Clonality :

Monoclonal

Clone No. :

TSKL10088

Reactivity :

Human,Mouse,Rat

Storage :

-15°C to -25°C/1 year
(Do not lower than -25°C)

Specificity :

Endogenous

Purification :

Protein A

Molecular Weight :

MW(Calculated): 44 kDa

MW(Observed): 53 kDa

Application :

WB,IF,IP,ELISA

Dilution :

WB 1:1000-1:4000

IF 1:50-1:200

ELISA 1:5000-1:20000


IP 1:50-1:200

Storage Buffer :

PBS,50%glycerol,0.05%Proclin
300,0.05%BSA

Western blot analysis of various lysates using p53(Phospho Ser392) Rabbit mAb (TS10088) at 1:2000 dilution.

Image



Lane 1: HEK293 cell lysate

Lane 2: Etoposide-treated HEK293 cell lysate

Lane 3: NIH-3T3 cell lysate

Lane 4: Etoposide-treated NIH-3T3 cell lysate

Lysates/proteins at 20 μg/Lane

Predicted band size: 44 kDa

Observed band size: 53 kDa

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5%BSA in TBST.

Exposure time: Lane 1-2:10 s(pg-level);Lane 3-4:25 s (fg-level).

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster

Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;

IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation

01

p53(Phospho Ser392) Rabbit mAb

Catalog :

TS10088

Gene Name :

p53

Gene ID :

Human: 7157

Mouse: 22059

Rat: 24842

SwissProt :

Human: P04637

Mouse: P02340

Rat: P10361

Host :

Rabbit

Clonality :

Monoclonal

Clone No. :

TSKL10088

Reactivity :

Human,Mouse,Rat

Storage :

-15°C to -25°C/1 year
(Do not lower than -25°C)

Specificity :

Endogenous

Purification :

Protein A

Molecular Weight :

MW(Calculated): 44 kDa

MW(Observed): 53 kDa

Application :

WB,IF,IP,ELISA

Dilution :

WB 1:1000-1:4000

IF 1:50-1:200

ELISA 1:5000-1:20000


IP 1:50-1:200

Storage Buffer :

PBS,50%glycerol,0.05%Proclin
300,0.05%BSA

Western blot analysis of various lysates using p53 (Phospho Ser392) Rabbit mAb (TS10088) at 1:2000 dilution.

Image



Lane 1: NIH-3T3 cell lysate

Lane 2: Staurosporine-treated NIH-3T3 cell lysate

Lane 3: C6 cell lysate

Lane 4: Staurosporine-treated C6 cell lysate

Lysates/proteins at 20 μg/Lane

Predicted band size: 44 kDa

Observed band size: 53 kDa

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG(H+L) at 1:10000 dilution. Blocking buffer: 5%BSA in TBST.

Exposure time: 16 s(fg-level).

Reactivity Key: H=Human;M=Mouse;R=Rat;Mk=Monkey;Hm=Hamster

Application Key:WB=Western Blot;IHC-P=Immunohistochemistry(Paraffin);ICC=Immunocytochemistry;

IF=Immunofluorescence;FC=Flow Cytometry;IP=Immunoprecipitation

02